

ALGEBRAIC MONOIDS AND GROUP EMBEDDINGS

ALVARO RITTATORE

ABSTRACT. We study the geometry of algebraic monoids. We prove that the group of invertible elements of an irreducible algebraic monoid is an algebraic group, open in the monoid. Moreover, if this group is reductive, then the monoid is affine. We then give a combinatorial classification of reductive monoids by means of the theory of spherical varieties.

1. INTRODUCTION

Let k be an algebraically closed field of arbitrary characteristic. An *algebraic monoid* is an algebraic variety S over k with an associative product $m : S \times S \rightarrow S$ which is a morphism of varieties, and such that there exists an element $1 \in S$ with $m(1, s) = m(s, 1) = s$ for all $s \in S$. We write $m(s, s') = ss'$ for all $s, s' \in S$. Let $G(S)$ be the *unit group* of S :

$$G(S) = \{g \in S : \exists g' \in S, gg' = g'g = 1\}.$$

We have the following picture:

$$\begin{array}{ccc} \{ \text{algebraic monoids} \} & \supset & \{ \text{algebraic groups} \} \\ \cup & & \cup \\ \{ \text{affine alg. monoids} \} & \supset & \{ \text{affine alg. groups} \} \end{array}$$

It is known that if S is an affine algebraic monoid, then $G(S)$ is an algebraic group, open in S ([8]). We extend this result to irreducible algebraic monoids. Moreover, the natural action of $G(S) \times G(S)$ on S by left and right multiplication has only one closed orbit : S is a *simple embedding* of $G(S)$. Furthermore, if we require $G(S)$ to be a reductive group, then the monoid S is affine. The proof of this result is based on the following remark:

If S is a reductive monoid, then by the Bruhat decomposition there exist opposite Borel subgroups B, B^- of $G(S)$ such that BB^- is open in S . So S is a spherical $(G(S) \times G(S))$ -variety. It follows that if S has a zero, then S is affine. We reduce the general case to that one by using techniques of the theory of spherical varieties.

Moreover, using the dictionary between simple spherical varieties and a family of combinatorial objects, the *colored cones* ([5]), we classify reductive monoids in arbitrary characteristic, in a way dual to that one of [11] (in characteristic zero).

Date: February 1, 2008.

Partially supported by a CONICYT's grant and the Universidad de la República (Uruguay).

I wish to thank M. Brion, L. Renner and E. Vinberg for many useful remarks and sugestions.

2. ALGEBRAIC MONOIDS AND GROUP EMBEDDINGS

We say that an algebraic monoid S is *irreducible* (resp. *normal*) if its underlying variety is irreducible (resp. normal).

An element $e \in S$ is an *idempotent (element)* of S if $e^2 = e$. We note $E(S)$ the set of idempotents of S .

We say that a monoid S has a *zero* if there exists an element $0 \in S$ such that $0s = s0 = 0$ for all $s \in S$.

An *ideal* of S is a non-empty subset $I \subset S$ such that $SIS \subset I$.

If there exists an ideal $Y \subset S$ contained in all the ideals of S , we say that S has *kernel* Y .

If the unit group of a monoid is a reductive algebraic group, we say that the monoid is *reductive*.

A morphism $\pi : S \rightarrow S'$ between two algebraic monoids is a *morphism of algebraic monoids* if $\pi(ss') = \pi(s)\pi(s')$, and $\pi(1_S) = 1_{S'}$. It is clear that $\pi(G(S)) \subset G(S')$. If S and S' have a zero, we impose that $\pi(0_S) = 0_{S'}$.

These conditions are not automatically satisfied:

Example 1. Let $M(n)$ be the monoid of $n \times n$ matrices with coefficients in k . We can identify $M(n)$ with the monoid of endomorphisms of a k -vector space V of dimension n . Its unit group is $\text{Gl}(n)$, the group of invertible matrices; it is isomorphic to the group of automorphisms of V . The monoid $M(n)$ is an affine, smooth, irreducible, reductive monoid with zero.

The morphism $M(n) \rightarrow M(n)$, $A \mapsto 0$, commutes with the multiplication, but it is not a morphism of algebraic monoids. The morphism $M(n) \rightarrow M(n) \times M(n)$, $A \mapsto (A, \text{id})$, is not a morphism of algebraic monoids. Indeed, the image of the zero matrix is not the zero of $M(n) \times M(n)$.

Notation. From now on we suppose that all the algebraic monoids are irreducible, unless otherwise mentioned.

Definition 1. Let G be an algebraic group. Consider the action of $G \times G$ on G given by

$$(a, b) \cdot g = agb^{-1} \quad , \quad a, b, g \in G .$$

This action is transitive, and the isotropy group of 1 is $\Delta(G)$, the diagonal of G . Moreover, the action satisfies the equation:

$$(1) \quad (a, 1) \cdot b = ab = (1, b^{-1}) \cdot a .$$

We say that a $(G \times G)$ -variety X is an *embedding of G* if there exists an element $x \in X$ such that the orbit $\mathcal{O}_x$ is open in X and isomorphic to $G \cong (G \times G)/\Delta(G)$. If there exists only one closed orbit, we say that X is a *simple embedding*.

Theorem 1. *Let S be an algebraic monoid. Then $G(S)$ is a connected algebraic group, open in S , and S is a simple embedding of $G(S)$. Moreover, the unique closed $(G(S) \times G(S))$ -orbit in S is the kernel of S .*

PROOF. Let $m : S \times S \rightarrow S$ be the product in S , and $p_1, p_2 : S \times S \rightarrow S$ the projections. Then

$$(2) \quad G = p_1(m^{-1}(1)) \cap p_2(m^{-1}(1)) ,$$

and G is constructible in S . If we prove that G is open and dense in S , it will follow in particular that G is connected, because S is irreducible.

From equation (2), we have

$$G \supset p_1(m^{-1}(1)^\circ) \cap p_2(m^{-1}(1)^\circ) ,$$

where $m^{-1}(1)^\circ$ is an irreducible component of $m^{-1}(1)$ that contains $(1, 1)$.

The product m is surjective, and S is irreducible, so ([4, prop. 6.4.5])

$$\dim m^{-1}(1)^\circ \geq \dim S .$$

Let $\rho_i = p_i|_{m^{-1}(1)^\circ}$, $i = 1, 2$. In order to show that G is dense in S , it is sufficient to prove that the morphisms ρ_i are dominant: then there exists a non-empty open subset of S contained in $\rho_1(m^{-1}(1)^\circ) \cap \rho_2(m^{-1}(1)^\circ) \subset G$.

The fibre $\rho_1^{-1}(1)$ is composed of the couples $(x, y) \in S \times S$ such that $x = 1$ and $xy = 1$, so $\rho_1^{-1}(1) = \{(1, 1)\}$.

Because of the upper-semicontinuity of the function $x \mapsto \dim \rho^{-1}(\rho(x))$, there exists a non-empty open subset $U \subset m^{-1}(1)^\circ$ such that $\dim \rho^{-1}(\rho(x)) = 0$ for all $x \in U$. Therefore,

$$\dim S \leq \dim m^{-1}(1)^\circ = \dim \overline{\rho_1(m^{-1}(1)^\circ)} \leq \dim S .$$

We have then equality, and $\dim \overline{\rho_1(m^{-1}(1)^\circ)} = \dim S$, and thus ρ_1 is a dominant morphism. By symmetry, ρ_2 is a dominant morphism too.

The left multiplication $\ell_g : G \times S \rightarrow S$ by an element $g \in G$ is an automorphism of S , with inverse $\ell_{g^{-1}}$, such that $\ell_g(G) = G$. It follows immediately that G is open in S , and that every point in G is smooth in S .

Next, we show that $g \mapsto g^{-1}$ is a morphism. Before doing that, we prove that if $s \in S$ has a left inverse, then s is a unit.

Let $s' \in S$ be such that $s's = 1$. Then ℓ_s has $\ell_{s'}$ as a left inverse, so ℓ_s is injective. The group G being open in S , $\ell_s(G)$ is constructible in S and $\dim G = \dim S$. Then $\ell_s(G)$ contains a non-empty open subset U of S . But $U \cap G \neq \emptyset$,

so $\ell_s(G) \cap G \neq \emptyset$. If $g \in G \cap \ell_s(G)$, there exists $g' \in G$ such that $g = sg'$, and thus $s = gg'^{-1} \in G$. Therefore, we have:

$$m^{-1}(1) = \{(x, y) \in S \times S : x \in G, y = x^{-1}\},$$

and $m^{-1}(1) = m^{-1}(1)^\circ$ is irreducible, so that $p_1|_{m^{-1}(1)} = \rho_1 : m^{-1}(1) \rightarrow G$ is bijective. If ρ_1 is separable, then it follows from Zariski's Main Theorem that ρ_1 is an isomorphism. Analogously, ρ_2 is an isomorphism, and the inverse map $g \mapsto g^{-1}$ is obtained by the composition $\rho_2 \circ \rho_1$, so it is an isomorphism.

So we must prove that ρ_1 is separable. Every point of G is smooth in S , so we can consider the differential of the product m at the point $(1, 1) \in G \times G$:

$$T_{(1,1)}m : T_{(1,1)}(S \times S) \cong T_1S \times T_1S \rightarrow T_1S \quad , \quad (u, v) \mapsto u + v.$$

Indeed, $m|_{\{1\} \times S}(s) = s$ and $m|_{S \times \{1\}}(s) = s$ for all $s \in S$; the restriction of $T_{(1,1)}m$ to the subspaces $T_1S \times \{0\}$ and $\{0\} \times T_1S$ is then the identity.

In particular, $T_{(1,1)}m$ is surjective, and the subvariety $m^{-1}(1) \subset S \times S$ is smooth at $(1, 1)$, with tangent space equal to

$$T_{(1,1)}m^{-1}(1) = \ker T_{(1,1)}m = \{(u, v) \in T_1S \times T_1S : u + v = 0\}.$$

The differential of ρ_1 at the point $(1, 1)$ is the restriction to $T_{(1,1)}m^{-1}(1)$ of the differential of the projection. It follows that $T_{(1,1)}\rho_1$ is surjective, and ρ_1 is separable.

We now show that S is a simple embedding of $G = G(S)$. We consider the action of $G \times G$ over S by left and right multiplication :

$$(g_1, g_2) \cdot s = g_1 s g_2^{-1} \quad , \quad g_1, g_2 \in G, s \in S.$$

The $(G \times G)$ -orbit passing by 1 is G , so S is an embedding of G . If Y is a closed $(G \times G)$ -orbit, then

$$SYS = \overline{GYG} \subset \overline{GYG} = \overline{Y} = Y.$$

It follows that every closed orbit is an ideal. If Y and Y' are closed orbits, then $\emptyset \neq YY' \subset Y \cap Y'$, and thus $Y = Y'$, because Y and Y' are orbits. Let Y be the unique closed orbit. If $I \subset S$ is an ideal, then $\emptyset \neq IY \subset Y \cap I$, and it follows that $Y \subset I$, so Y is the kernel of S . $\square$

We remark that in the proof of the preceding theorem we have proved :

Corollary 1. *Let S be an algebraic monoid. If $s \in S$ has a left (or right) inverse, then s is an unit.* $\square$

On the other side, we have a partial converse to theorem 1 (which was proved by Vinberg in characteristic 0 – see [11]):

Proposition 1. *Let G be an algebraic group and S be an affine embedding of G . Then the product on G extends to a product $\tilde{m} : S \times S \rightarrow S$, in such a way that $G(S) = G$.*

PROOF. If the product $m : G \times G \rightarrow G$ extends to a morphism $\tilde{m} : S \times S \rightarrow S$, the $\tilde{m}$ is a product, because $\tilde{m}|_{G \times G} = m$ is associative. For convenience, we note $\tilde{m} = m$. It is clear that the identity $1 \in G$ satisfies $1s = s1 = s$ for all $s \in S$. Indeed, this holds for the elements of the open subset $G \subset S$. Consider the right and left actions of G given by :

$$\begin{aligned} G \times S &\rightarrow S & g \cdot s &= (g, e) \cdot s \\ S \times G &\rightarrow S & s \cdot g &= (e, g^{-1}) \cdot s \end{aligned}$$

By construction and equation (1), these actions coincide on $(G \times G) \subset (S \times S)$ with the product in G . Then there exists a morphism

$$m : (G \times S) \cup (S \times G) = U \rightarrow S$$

which extends the product of G . Therefore, it suffices to prove that every regular function defined in U can be extended to $S \times S$.

If V is a vector space and $W \subset V$ a subspace, an easy calculation shows that $W \otimes W = (V \otimes W) \cap (W \otimes V)$. Applying this to $k[S] \subset k[G]$, we get:

$$k[S \times S] \cong k[S] \otimes k[S] = (k[S] \otimes k[G]) \cap (k[G] \otimes k[S]) .$$

On the other hand, a function $f \in k[G] \otimes k[G]$ belongs to $k[S] \otimes k[G]$ if and only if it can be extended to a function defined in $S \times G$. The same happens for $k[G] \otimes k[S]$. It follows that every function defined in U can be extended to a function defined in $S \times S$.

Finally, we prove that $G(S) = G$. The inclusion $G \subset G(S)$ is trivial. Let $x \in S \setminus G$ be a unit. Then $\ell_x(x^{-1}) = x \cdot x^{-1} = 1 \in G$, and there exists $g \in G$ such that $x \cdot g \in G$, because G is open in S . It follows that $x = (x \cdot g) \cdot g^{-1} \in G$. $\square$

In order to study the geometry of monoids it is useful to suppose that they are normal. The following result allows us to doing so without loss of generality:

Lemma 1. ([9], [10]) *Let S be an algebraic monoid, and $\pi : \tilde{S} \rightarrow S$ its normalization. Then there exists an unique associative product $\tilde{m} : \tilde{S} \times \tilde{S} \rightarrow \tilde{S}$ such that $\pi \circ \tilde{m} = m \circ (\pi, \pi)$. Moreover, the following proprieties are verified:*

- i) $I \subset S$ is an ideal if and only if $\pi^{-1}(I) \subset \tilde{S}$ is an ideal.
- ii) S has a zero 0_S if and only if $\pi^{-1}(0_S) = \{0_{\tilde{S}}\}$, where $0_{\tilde{S}}$ is a zero of $\tilde{S}$.
- iii) $\tilde{S}$ is an algebraic monoid, with $G(\tilde{S}) = \pi^{-1}(G(S))$, and

$$\pi|_{G(\tilde{S})} : G(\tilde{S}) \rightarrow G(S)$$

is an isomorphism of groups.

- iv) The restriction of π to $E(\tilde{S})$ gives a bijection between $E(\tilde{S})$ and $E(S)$.

It follows that π is a morphism of (algebraic) monoids. In particular, π is equivariant for the actions of $G(\tilde{S}) \times G(\tilde{S})$ and $G(S) \times G(S)$.

PROOF. We define the multiplication $\tilde{m} : \tilde{S} \times \tilde{S} \rightarrow \tilde{S}$ by means of the universal propriety of normalizations: $\tilde{m}$ is the unique morphism such that the diagram

$$\begin{array}{ccc} \tilde{S} \times \tilde{S} & \xrightarrow{\tilde{m}} & \tilde{S} \\ \pi \times \pi \downarrow & & \downarrow \pi \\ S \times S & \xrightarrow{m} & S \end{array}$$

is commutative.

By the universal property of normalizations, we can easily prove that $\tilde{m}$ is associative, and by construction, π is then a morphism of algebraic semigroups.

We prove only iv). For the others assertions, we refer the reader to [10].

It is clear that $\pi(E(\tilde{S})) \subset E(S)$. Let $e \in E(S)$, and $e_1 \in \pi^{-1}(e)$. The morphisms $\ell_{e_1 e_1} = \ell_{e_1} \ell_{e_1}$ and ℓ_{e_1} verify the following commutative diagram:

$$\begin{array}{ccccc} \tilde{S} & \xrightarrow{\tilde{\ell}_{e_1 e_1}} & \tilde{S} & \xleftarrow{\tilde{\ell}_{e_1}} & \tilde{S} \\ \pi \downarrow & & \downarrow \pi & & \downarrow \pi \\ S & \xrightarrow{\ell_e \ell_e = \ell_e} & S & \xleftarrow{\ell_e} & S \end{array}$$

Again by the universal property of normalizations, $\tilde{\ell}_{e_1} = \tilde{\ell}_{e_1 e_1}$. In particular, $e_1 = \tilde{\ell}_{e_1}(1) = \tilde{\ell}_{e_1 e_1}(1) = e_1 e_1$, so $\pi^{-1}(E(S)) \subset E(\tilde{S})$.

Let $e_1, e_2 \in \pi^{-1}(e)$. If we replace $\tilde{\ell}_{e_1 e_1}$ by $\tilde{\ell}_{e_1 e_2}$ in the preceding diagram, we get again a commutative diagram, and it follows that $e_1 e_2 = e_1$. Analogously, $e_1 e_2 = e_2$, and thus $e_1 = e_2$. $\square$

3. REDUCTIVE MONOIDS

We restrict ourselves to the study of reductive (irreducible) monoids. All the reductive groups are supposed connected.

Definition 2. If G is a reductive group, a homogeneous space G/H is *spherical* if there exists a Borel subgroup B of G such that $BH \subset G$ is open. If X is a G -variety with an open orbit isomorphic to G/H , then X is called an *embedding of G/H* . We say that X is *simple* if the action of G over X has only one closed orbit. If X is normal, we say that X is a *spherical variety*.

If G is a reductive group, it follows from the Bruhat decomposition that the normal G -embeddings are spherical varieties, so, by Theorem 1, normal reductive monoids are simple spherical varieties. The aim of this section is to study the geometry of reductive monoids from the point of view of the theory of spherical varieties. In particular, we will show that reductive monoids are affine.

Remark. In the litterature, it is imposed to G -embeddings to be normal. We don't do this in order to be able to state results valid for all reductive monoids.

We begin by proving the following

Lemma 2. *Let S be a reductive monoid with zero. Then S is affine.*

PROOF. More generally, if X is a simple spherical variety, with closed orbit a point, then X is affine ([5, thms. 7.6 and 7.7]). If S is not normal, it suffices to take the normalization $\pi : \tilde{S} \rightarrow S$. It follows from Lemma 1 that $\tilde{S}$ is a normal monoid with zero. So $\tilde{S}$ is affine, and it follows that S is. Indeed, the morphism of the normalization π is a finite surjective morphism, so $\tilde{S}$ is affine if and only if S is ([2, p. 63]). $\square$

In order to continue our study, we must give *a priori* proofs of results concerning idempotents in reductive monoids, which are known in the affine case. We recall that an *1 parameter subgroup* (1-PS) of a group G is a multiplicative morphism $\lambda : k^* \rightarrow G$. We note $\Xi_*(G)$ the group of 1-PS of G .

Lemma 3. *Let S be a reductive monoid with unit group G , and let $T \subset G$ be a maximal torus. Then every $(G \times G)$ -orbit contains an idempotent element belonging to $\overline{T}$.*

PROOF. First, we show that we can approach every $(G \times G)$ -orbit of S by an 1-PS of G .

If $x \in S$, there exists a curve germ $\xi \in G_{((t))} = \text{Hom}(\text{Spec } k((t)), G)$, where $k((t))$ denotes the ring of Laurent series in one variable with coefficients in k , such that $\lim_{t \rightarrow 0} \xi(t) = x$. By Iwahori's Theorem ([6]), there exists an 1-PS λ of G such that $\lambda = f\xi f'$, where f and f' belong to $G_{k[[t]]} = \text{Hom}(\text{Spec } k[[t]], G)$. Let $g = f(0)$ and $g' = f'(0)$. Then,

$$\lim_{t \rightarrow 0} \lambda(t) = \lim_{t \rightarrow 0} f\xi(t)f' = g x g' \in \mathcal{O}_x .$$

The 1-PS λ has its values in a maximal torus T' . As all maximal tori of G are conjugated, there exists $g_0 \in G$ such that $g_0 \lambda(t) g_0^{-1} \in T$ for all $t \in k^*$. Then $\lambda_0(t) = g_0 \lambda(t) g_0^{-1} \in \Xi_*(T)$ has a limit in $\mathcal{O}_x$.

Next, we observe that if $\lim_{t \rightarrow 0} \lambda_0(t) = e$, then

$$e^2 = \lim_{t \rightarrow 0} \lambda(t) \lim_{t \rightarrow 0} \lambda(t) = \lim_{t \rightarrow 0} \lambda(t)^2 = \lim_{t \rightarrow 0} \lambda(t^2) = e ,$$

so e is the idempotent we are looking for. $\square$

Lemma 4. *Let S be a reductive monoid with unit group G . Let $e \in E(S)$, and $Y = \mathcal{O}_e = GeG$. If Y is a monoid with identity e , then e is a central element of S , and Y is an algebraic group. The isotropy group of e satisfies:*

$$(G \times G)_e = (G_1 \times G_1) \Delta(G) ,$$

where G_1 is the isotropy group of e for the action of G by left multiplication. Moreover, G_1 is a normal subgroup of G . In particular, $(G \times G)_e$ is a reductive group.

PROOF. As e is the identity of Y , it follows that $ge = e(ge) = (eg)e = eg$ for all $g \in G$. Then $(g, g) \cdot e = geg^{-1} = gg^{-1}e = e$, and $\Delta(G) \subset (G \times G)_e$. Moreover, $G \subset S$ is dense in S , so $se = es$ for all $s \in S$. Thus, e is a central element of S .

In particular $Y = GeG = Ge$, and $\varphi_0 : G/G_1 \rightarrow Y$ is a bijective morphism. Observe that the multiplication in Y has an inverse, namely $(ge)^{-1} = g^{-1}e$.

If $g_1 \in G_1$ and $g \in G$, then

$$(gg_1g^{-1}) \cdot e = gg_1g^{-1}e = gg_1eg^{-1} = geg^{-1} = gg^{-1}e = e.$$

It follows that G_1 is normal in G . Thus, φ_0 is compatible with products.

On the other hand, it is clear that $(G_1 \times G_1)\Delta(G) \subset (G \times G)_e$. If $(a, b) \in (G \times G)_e$, then

$$e = (a, b) \cdot e = aeb^{-1} = ab^{-1}e = (ab^{-1}, 1) \cdot e.$$

It follows that $ab^{-1} \in G_1$, and $(a, b) = (ab^{-1}, 1)(b, b) \in (G_1 \times G_1)\Delta(G)$. $\square$

Remark. The multiplication in Y may not be induced by that one of $G \times G$ through $G \times G \rightarrow (G \times G)/(G \times G)_e \rightarrow Y$. Indeed, $(G \times G)_e$ is not always normal in $G \times G$.

The following result of Waterhouse shows some properties of unit groups of affine algebraic monoids.

Proposition 2. ([13],[9, thm. 3.3.6]) *Let G be an algebraic group (not necessarily reductive). Then the following properties are equivalent:*

- i) *The character group of G is non trivial.*
- ii) *The rank of the radical of G is not equal to zero.*
- iii) *There exists an affine algebraic monoid with unit group G and such that $G \neq S$.*

If moreover G is reductive, we show that the *connected center* of S (i.e. the closure in S of Z , the connected center of G) meets the kernel of S :

Proposition 3. *Let S be a reductive monoid with unit group $G \neq S$; let Z be the connected center of G , and Y be the kernel of S . Then $\overline{Z} \cap Y \neq \emptyset$. In particular, the connected center of G is non trivial.*

PROOF. Let T be a maximal torus of G . By Lemma 3, there exists an 1-PS $\lambda \in \Xi_*(T)$ whose limit exists and is an idempotent $e \in Y \cap \overline{T}$.

If $W = N_G(T)/C_G(T)$ is the Weyl group of G , then W acts on T by conjugation. We obtain then an action of W over $\Xi_*(T)$ as follows:

$$w \cdot (\mu(t)) = w\mu(t)w^{-1} \in \Xi_*(T) \quad \forall \mu \in \Xi_*(T), w \in W.$$

From $W \cdot T \subset T$ we obtain $W \cdot \overline{T} \subset \overline{T}$. Let $\mu = \sum_{w \in W} w \cdot \lambda \in \Xi_*(T)$; it is a W -invariant 1-PS, with limit $e = \prod_{w \in W} wew^{-1} \in E(Y)$. In this way, we have constructed a 1-PS μ of the center of G (because it is W -invariant) whose limit e belongs to Y . In particular, μ is non trivial, and $e \in Y \cap \overline{Z}$: the proposition is proved. $\square$

It is known that every affine embedding of a semi-simple group is trivial ([8]). This result follows immediately from the proposition above:

Corollary 2. *If the unit group of a monoid S is a semi-simple group G_0 , then $G_0 = S$.* $\square$

Theorem 2. *Let S be a reductive monoid. Then S is affine.*

PROOF. First, we consider the case where S is normal. Consider the kernel Y of S , $G = G(S)$, $T \subset G$ a maximal torus and $Z \subset G$ the connected center. Let $e \in \overline{Z} \cap Y$ as in Proposition 3. Then e is a central element of S , so $es = se$ for all $s \in S$. An element $y \in Y$ can be written as a product $y = geg' = gg'e = egg'$. In particular, Y is an algebraic monoid with identity $e \in Y$.

Consider the morphism $\psi : S \rightarrow Y$, $s \mapsto se$. We have:

$$\begin{cases} \psi((g, g') \cdot s) = \psi(gsg'^{-1}) = gsg'^{-1}e = gseg'^{-1} = (g, g') \cdot (se) = (g, g') \cdot \psi(s) \\ \psi(ss') = ss'e = ss'ee = ses'e = \psi(s)\psi(s') \end{cases}$$

It follows that ψ is an $(G \times G)$ -equivariant morphism of monoids. The fibre $F_e = \{s \in S \mid se = e\}$ is then stable under multiplication and under the action of $(G \times G)_e$.

The identity 1 belongs to F_e : the fibre F_e is a monoid with zero (equal to e). If $G_1 = \{g \in G \mid ge = e\}$, from Lemma 4 we deduce that $(G \times G)_e = (G_1 \times G_1)\Delta(G)$. Moreover, $F_e \cap G = G_1$, and it follows that G_1 is an open subset of F_e , so $G_1 = G(F_e)$ by Theorem 1. Then F_e is a reductive monoid with zero, so it is affine.

We consider the affine variety $F = (G \times G) \times F_e$. The reductive group $(G \times G)_e$ acts in F :

$$(a, b) \cdot ((g_1, g_2), s) = ((g_1 a^{-1}, g_2 b^{-1}), asb^{-1}),$$

where $(a, b) \in (G \times G)_e$, $g_1, g_2 \in G$, $s \in F_e$. The quotient $(G \times G)_{*(G \times G)_e} F_e = F // (G \times G)_e$ is then an affine variety. If we prove that this quotient is isomorphic to S , it will follow that S is affine.

We consider the morphism

$$\gamma : F = (G \times G) \times F_e \rightarrow S, \quad \gamma((a, b), s) = (a, b) \cdot s = asb^{-1}.$$

As γ is constant over the $(G \times G)_e$ -orbits of F , then γ induces a morphism $\tilde{\gamma} : (G \times G) *_{(G \times G)_e} F_e \rightarrow S$. We show that $\tilde{\gamma}$ is bijective and birational, then it follows from Zariski's Main Theorem that $\tilde{\gamma}$ is an isomorphism.

$\tilde{\gamma}$ is injective : Let $((a, b), s), ((a', b'), s') \in (G \times G) \times F_e$ such that $\gamma((a, b), s) = \gamma((a', b'), s')$. Then $(a, b) \cdot s = (a', b') \cdot s'$, and $s' = (a'^{-1}a, b'^{-1}b) \cdot s$. It follows that

$$(a, b) \cdot e = \psi((a, b) \cdot s) = \psi((a', b') \cdot s') = (a', b') \cdot e.$$

Then $(a'^{-1}a, b'^{-1}b) \cdot e = e$, and thus $(a'^{-1}a, b'^{-1}b) \in (G \times G)_e$. So

$$\overline{((a', b'), s')} = \overline{((a', b'), (a'^{-1}a, b'^{-1}b) \cdot s)} = \overline{((a, b), s)}.$$

$\tilde{\gamma}$ is surjective : If $s \in S$, then $\psi(s) = se = ge$ for some $g \in G$. Then $((g, 1), g^{-1}s) \in (G \times G) \times F_e$, and $\gamma((g, 1), g^{-1}s) = s$.

$\tilde{\gamma}$ is birational :

The group $G \times G$ acts on $(G \times G) *_{(G \times G)_e} F_e$ by left multiplication in the left factor, in such a way that $\tilde{\gamma}$ is $(G \times G)$ -equivariant. Moreover, the orbit $\mathcal{O}_{((1,1),1)}$ passing by $((1, 1), 1)$ is open in $(G \times G) *_{(G \times G)_e} F_e$, and isomorphic to the quotient $(G \times G) *_{(G \times G)_e} G_1$. The restriction of $\tilde{\gamma}$ gives then an isomorphism from $\mathcal{O}_{((1,1),1)}$ to G , with inverse $g \mapsto \overline{((g, 1), 1)} \in (G \times G) *_{(G \times G)_e} G_1$.

For the general case, observe that if S reductive, non-normal monoid, its normalization $\tilde{S}$ is a reductive normal monoid by Lemma 1. It follows that $\tilde{S}$, and therefore S , is affine. Indeed, the normalization $\pi : \tilde{S} \rightarrow S$ is a finite surjective morphism.

□

Corollary 3. *The kernel of a normal reductive monoid is a reductive group.*

PROOF. We have seen that if S is a normal reductive monoid and Y its kernel, then Y is a group, with identity $e \in \overline{Z} \cap Y$. Moreover, if we keep the notation of the proof of theorem 2, then $S \cong (G \times G) *_{(G \times G)_e} F_e$. It follows that

$$Y \cong (G \times G) \cdot \overline{((1, 1), e)} \cong (G \times G) / (G \times G)_e.$$

But $(G \times G) / (G \times G)_e \cong G / G_1$, and Y is a reductive group.

□

Remarks. i) Taking irreducible components, it is easy to prove that if S is a (reducible) reductive normal monoid, then $\overline{G(S)}$ is affine, and each unit belongs to a unique irreducible component ([10]). Moreover, any two irreducible components of $\overline{G(S)}$ are isomorphic as varieties.

ii) It is known that any irreducible affine monoid can be embedded as a closed submonoid of the monoid of endomorphisms of some vector space V of finite dimension ([8]). It follows from Theorem 2 that any reductive monoid S can be

seen as the closure in $\text{End}(V)$ of a closed reductive subgroup of $\text{Gl}(V)$, which is then isomorphic to $G(S)$.

We can summarize our results in the following

Theorem 3. *The reductive monoids are exactly the affine embeddings of reductive groups. The commutative reductive monoids are exactly the affine embeddings of tori.*

PROOF. We have only to prove the last statement. It is clear that if S is a affine embedding of a torus then S is commutative. Conversely, if S is a reductive commutative monoid, then its unit group is a reductive commutative connected group, i.e. a torus. $\square$

Let S be a reductive monoid, and G its unit group. If T is a maximal torus of G , then the properties of $\overline{T}$ reflect in some sense those of S . We state first a known result on irreducible monoids whose unit group is a torus (see for example ([7]).

Lemma 5. *Let S be a monoid whose unit group is a torus T . Then the ideals of S are exactly the T -stable closed subsets of S . In particular, every ideal of S is closed. Moreover, if $e, e' \in E(S)$ then $ee' = e'$ if and only if $e' \in \overline{T}e$.* $\square$

This result generalizes to reductive monoids as follows :

Proposition 4. *Let S be a reductive monoid, with unit group G . Then the ideals of S are exactly the $(G \times G)$ -stable closed subsets of S . In particular all ideals are closed.*

Two idempotents $e, e' \in E(S)$ are in the same $(G \times G)$ -orbit if and only if they are G -conjugate. In particular, if $T \subset G$ is a maximal torus, then

$$E(S) = \bigcup_{g \in G} gE(\overline{T})g^{-1}.$$

PROOF. We prove the first part. For the second one, see for example [8, cor. 6.10].

If $I \subset S$ is a $(G \times G)$ -stable closed subset, then

$$SIS = \overline{GIG} \subset \overline{GIG} = \overline{I} = I,$$

that is, I is an ideal of S .

Let $I \subset S$ be an ideal of S . By definition, I is $(G \times G)$ -stable. Let us prove that I is closed.

From lemma 5, we get that

$$I = \cup_{i=1}^n Ge_iG, \quad e_i \in I \cap \overline{T} \cap E(S).$$

Then, its closure in S is equal to $\cup_{i=1}^n \overline{Ge_iG}$. We need only to prove that if $e \in E(S) \cap I \cap \overline{T}$, then $\overline{GeG} \subset I$.

Note that $\overline{GeG} = \cup_{j=1}^m G\varepsilon_j G$, with $\varepsilon_j \in \overline{GeG} \cap \overline{T}$ an idempotent element. Moreover, $\overline{GeG}$ is a $(G \times G)$ -stable closed subset of S , so it is an ideal of S . The intersection $GeG \cap \overline{T}$ is a non-empty T -stable subset of $\overline{GeG} \cap \overline{T}$, so

$$GeG \cap \overline{T} = \cup_{i=1}^l Tg_i eg'_i T = \cup_{i=1}^l Tg_i eg'_i \quad g_i, g'_i \in G \quad , \quad g_i eg'_i \in E(S) \quad ,$$

and the closure of $GeG \cap \overline{T}$ is an ideal of $\overline{T}$. Let us call this ideal J . As $\varepsilon_j \in \overline{GeG} \cap \overline{T}$ for all $j = 1, \dots, m$, we get $\varepsilon_j e \in J \cap E(\overline{T})$, and there exists $i \in \{1, \dots, l\}$ such that $\varepsilon_j e \in Tg_i eg'_i$. It follows from lemma 5 that

$$\varepsilon_j e = \varepsilon_j eg_i eg'_i = (\varepsilon_j eg_i) eg'_i \in SeS \subset I \quad ,$$

and the proposition is proved. $\square$

4. CLASSIFICATION OF REDUCTIVE MONOIDS

In [11], Vinberg classified all reductive normal monoids in characteristic zero in terms of the decomposition of their algebra of regular functions for the action of the unit group. This classification is dual of that of affine embeddings of a reductive group (see [12]). From the work of Knop ([5]), we know how to classify affine embeddings of reductive group in arbitrary characteristic. In order to do that, we begin by recalling some results from the theory of spherical embeddings and in particular *reductive embeddings* (i.e. embeddings of reductive groups), which proofs can be found in [5]. Next, we classify (normal) reductive monoids. In view of Lemma 1, we suppose that all varieties considered are normal.

4.1. Reductive embeddings.

Notation. Let X be an algebraic variety. We denote by $k(X)$ the field of rational functions over X .

Definition 3. Let G be a reductive group, G/H a spherical homogenous space, and X a simple spherical embedding of G/H , with unique closed orbit Y . We note $\mathcal{D} = \mathcal{D}(G/H)$ the *set of colours*, i.e. the set of B -stable irreducible divisors of G/H . We call the *(set of) colours* of X , and we denote by $\mathcal{F}(X)$, the subset of $\mathcal{D}$ consisting of the irreducible divisors such that their closure contains Y .

We denote by $\mathcal{B}(X)$ the set of irreducible G -stable divisors of X .

Let $\Lambda_{G/H}$ be the set of weights of $k(G/H)^{(B)}$, we define

$$\mathcal{Q}(G/H) = \text{Hom}_{\mathbb{Z}}(\Lambda_{G/H}, \mathbb{Q}) \cong \text{Hom}_{\mathbb{Z}}(\Lambda_{G/H}, \mathbb{Z}) \otimes_{\mathbb{Z}} \mathbb{Q} \quad .$$

We call the *rank* of G/H the dimension of the $\mathbb{Q}$ -vector space $\mathcal{Q}(G/H)$.

Let M be an algebraic variety. A *valuation* of the field $k(X)$ is a function $\nu : k(X) \rightarrow \mathbb{Q} \cup \{-\infty\}$ with the following properties:

- i) $\nu(f_1 + f_2) \geq \min\{\nu(f_1), \nu(f_2)\}$ for all $f_1, f_2 \in k(X)$.
- ii) $\nu(f_1 f_2) = \nu(f_1) + \nu(f_2)$ for all $f_1, f_2 \in k(X)$
- iii) $\nu(k^*) = 0$, $\nu(0) = -\infty$.

It is easy to prove that $\nu(k^*) = 0$, $\nu(0) = -\infty$. We say that the valuation ν is *normalized* if $\nu(k(X) \setminus \{0\}) = \mathbb{Z}$.

If X is a G -variety, a valuation ν of $k(X)$ is called G -invariant if $\nu(g \cdot f) = \nu(f)$ for all $g \in G$, $f \in k(X)$, where the G -action over $k(X)$ is given by $(g \cdot f)(x) = f(g^{-1}x)$ for all $x \in X$.

Let G/H be an spherical homogeneous space, ν a valuation of $k(G/H)$ and $\chi \in \Lambda_{G/H}$ a weight. If $f_\chi, f'_\chi \in k(G/H)^{(B)}$ are two rational functions of weight χ , then the function f_χ/f'_χ is constant over the open B -orbit, and thus it is constant over G/H . It follows that $\nu(f_\chi) = \nu(f'_\chi)$. We set $\rho_\nu(\chi) = \nu(f_\chi)$. If f_χ and $f_{\chi'}$ are two rational functions of weight χ and χ' respectively, then $f_\chi f_{\chi'}$ is a rational function of weight $\chi + \chi'$, and $\rho_\nu(\chi + \chi') = \rho_\nu(\chi) + \rho_\nu(\chi')$. Thus, every valuation ν of $k(G/H)$ defines an element $\rho_\nu \in \mathcal{Q}(G/H)$.

If $\mathcal{V} = \mathcal{V}(G/H)$ is the set of G -invariant valuations of $k(G/H)$, the restriction of ρ to $\mathcal{V}$ gives an injection $\rho : \mathcal{V} \hookrightarrow \mathcal{Q}(G/H)$ ([5]). We will identify $\mathcal{V}$ with its image in $\mathcal{Q}(G/H)$.

The set $\mathcal{V}$ is a polyhedral cone (i.e. a cone generated by a finite number of elements of the lattice $\text{Hom}_{\mathbb{Z}}(\Lambda, \mathbb{Z})$), we call it the *valuation cone*.

Let $D \in \mathcal{B}(X)$ be a G -stable irreducible divisor of X , and ν_D the (normalized) G -invariant valuation associated to D . As D is determined by ν_D , the map

$$\rho : \mathcal{B}(X) \rightarrow \mathcal{V}(G/H) \quad , \quad D \mapsto \rho(\nu_D)$$

is injective. We will identify $\mathcal{B}(X)$ with its image by ρ .

On the other hand, there is a natural mapping $\mathcal{D}(G/H) \rightarrow \mathcal{Q}(G/H)$: to each color D we associate ρ_{ν_D} , where ν_D is the valuation of $k(G/H)$ associated to D . This mapping, noted ρ , is not necessarily injective ([5]).

Definition 4. Let G/H be a spherical homogenous space. A *colored cone* of $\mathcal{Q}(G/H)$ is a pair $(\mathcal{C}, \mathcal{F})$, where $\mathcal{F} \subset \mathcal{D}(G/H)$ is a subset such that:

- i) $\rho(\mathcal{F})$ does not contain 0,
- ii) $\mathcal{C} \subset \mathcal{Q}(G/H)$ is a polyhedral cone generated by $\rho(\mathcal{F})$, and a finite number of elements of $\mathcal{V}$,
- iii) $\text{int}(\mathcal{C}) \cap \mathcal{V} \neq \emptyset$, where $\text{int}(\mathcal{C})$ denotes the relative interior of $\mathcal{C}$.

A colored cone is *strictly convex* if $\mathcal{C}$ is.

A *colored face* of the colored cone $(\mathcal{C}, \mathcal{F})$ is a colored cone $(\tau, \mathcal{F}')$, with τ a face of $\mathcal{C}$, and $\mathcal{F}' = \mathcal{F} \cap \rho^{-1}(\tau)$.

Proposition 5. (see [5]) i) *There exists a bijection between the simple embeddings of G/H and the strictly convex colored cones of $\mathcal{Q}(G/H)$. This bijection*

associates to each variety X the colored cone $(\mathcal{C}(X), \mathcal{F}(X))$, where $\mathcal{C}(X)$ is the cone generated by $\mathcal{B}(X) \cup \rho(\mathcal{F}(X))$. We call $(\mathcal{C}(X), \mathcal{F}(X))$ the colored cone associated to X . We note $\mathcal{V}(X) = \mathcal{C}(X) \cap \mathcal{V}(G/H)$.

ii) The G -orbits of a simple spherical variety X are in bijection with the colored faces of its associated colored cone. This bijection is constructed as follows:

Let $Y \subset X$ be a G -orbit. We denote by $\mathcal{C}_Y(X)$ the cone generated by the image by ρ of the B -stable irreducible divisors which contain Y . If $\mathcal{F}(Y)$ is the set of colours such that their closure contains Y , then $(\mathcal{C}_Y(X), \mathcal{F}(Y))$ is the colored face we are looking for.

iii) If $P \subset G$ is the parabolic subgroup given by the intersection of the stabilizers of the colours in $\mathcal{D} \setminus \mathcal{F}(Y)$, then

$$\dim Y = \operatorname{rg}(G/H) - \dim \mathcal{C}_Y(X) + \dim G/P.$$

In particular, the unique closed orbit is a point if and only if the dimension of $\mathcal{C}(X)$ is equal to the rank of G/H , and the set of colours of the embedding is all $\mathcal{D}$.

iv) A spherical variety X is affine if and only if X is simple and there exists $\chi \in \Lambda_{G/H}$ such that

- (a) $\chi|_{\mathcal{V}(G/H)} \leq 0$
- (b) $\chi|_{\mathcal{C}(X)} = 0$
- (c) $\chi|_{\rho(\mathcal{D}(G/H) \setminus \mathcal{F}(X))} > 0$.

□

In [12], Vust established the combinatorial data associated to symmetric spaces, in particular to the homogenous space $G \cong (G \times G)/\Delta(G)$, where G is a reductive connected group, and $\operatorname{car} k = 0$. In order to carry out the classification of reductive monoids, we must generalize his results to arbitrary characteristic. This is essentially done in the literature, so we will state the results without proving them. For a proof, we refer the reader to [10].

Let us fix the notation:

We note G_0 the commutator of G . Thus, $G = ZG_0$, where $Z = Z(G)$ is the connected center of G . Moreover, $Z_0 = G_0 \cap Z$ is finite; G is then isomorphic to $(G_0 \times Z)/Z_0$:

$$k[G] \cong k[(G_0 \times Z)/Z_0] \cong (k[G_0] \otimes_k k[Z])^{Z_0}.$$

We fix T_0 , a maximal torus of G_0 , and $B_0 \subset G_0$, a Borel subgroup that contains T_0 . Let $\alpha_1, \dots, \alpha_l$ and $\omega_1, \dots, \omega_l$ be the simple roots and fundamental weights associated to (B_0, T_0) respectively. We set $T = T_0Z$, a maximal torus of G . We note W the Weyl group associated to (G, T) , and $C = C(G)$ the Weyl chamber of G associated to B .

If $B = B_0Z$, the weight lattice of G associated to (B, T) , satisfies:

$$\Xi(T) \cong (\Xi(T_0) \times \Xi(Z))^{Z_0} = \{(\lambda, \mu) \in \Xi(T_0) \times \Xi(Z) : \mu|_{Z_0} = \lambda|_{Z_0}\}.$$

It is a sub-lattice of finite index of $\Xi(G_0) \times \Xi(Z)$.

The reflection associated to a simple root α is noted as usual s_α .

From the Bruhat decomposition, it is easy to prove the following

Proposition 6. ([10]) *Let G be a reductive group. If we keep the preceding notations and identify $\text{Hom}_{\mathbb{Z}}(\Xi(T), \mathbb{Z})$ with $\Xi_*(T)$, then*

$$\begin{aligned} \mathcal{Q}((G \times G)/\Delta(G)) &= \mathcal{Q}(G) = \Xi_*(T) \otimes \mathbb{Q} \cong \\ &(\Xi_*(T_0) \otimes \mathbb{Q}) \oplus (\Xi_*(Z) \otimes \mathbb{Q}) = \mathcal{Q}(G_0) \oplus \mathcal{Q}(Z), \end{aligned}$$

where the lattice considered is $(\Xi_*(T_0) \times \Xi_*(Z))^{Z_0} \subset \Xi_*(T_0) \times \Xi_*(Z)$.

We identify $\Xi_*(T)$ with $\Xi(T)$ by means of a W -invariant form $\langle \cdot, \cdot \rangle$, in such a way that $\langle \omega_i, \alpha_j^\vee \rangle = \delta_{ij}$. Under this identification, $\mathcal{Q}(G) = \Xi(T) \otimes \mathbb{Q}$, $\mathcal{Q}(G)^* = \Xi(T) \otimes \mathbb{Q}$, and the valuation cone $\mathcal{V}(G) \subset \mathcal{Q}(G)$ is opposite to $C(G)$.

Moreover, the colours of G are the divisors $D_i = \overline{Bs_{\alpha_i}B^-}$, $i = 1, \dots, l$, with associated valuations $(\alpha_i^\vee, 0) \in (\Xi(G_0) \times \Xi(Z))^{Z_0}$. $\square$

It follows immediately that the behaviour under quotients by a finite central subgroup of the combinatorial data associated to reductive monoids is “good”:

Proposition 7. ([10],[11]) *Let G be a reductive group and $\Gamma \subset G$ a finite central subgroup. Then*

- i) $\mathcal{Q}(G/\Gamma) = \mathcal{Q}(G)$, where we consider the lattice $\Xi_*(T/\Gamma) = \Xi_*(T)^\Gamma \subset \Xi_*(T)$.
- ii) If $\pi : G \rightarrow G/\Gamma$ is the canonical projection, then π induces a bijection between $\mathcal{D}(G)$ and $\mathcal{D}(G/\Gamma)$.
- iii) Under the identification i), $\rho_G(\mathcal{D}(G)) = \rho_{G/\Gamma}(\mathcal{D}(G/\Gamma))$, and the valuation cones $\mathcal{V}(G)$ and $\mathcal{V}(G/\Gamma)$ coincide. In particular, π_* is the identity.
- iv) If S is a monoid with unit group G , then S/Γ is a reductive monoid, with unit group G/Γ . Moreover, if we denote $\pi : S \rightarrow S/\Gamma$ the quotient, then $\pi_* : \mathcal{Q}(G) \rightarrow \mathcal{Q}(G')$ is the identity, and under this identification we have $\mathcal{C}(S/\Gamma) = \mathcal{C}(S)$.

$\square$

Remark. In the case of reductive embeddings, the application $\rho : \mathcal{D} \rightarrow \mathcal{Q}(G)$ is injective. We identify then $\mathcal{D}$ with its image by ρ .

From this, we can recover the fact that a reductive monoid is a quasi-direct product of a reductive group and a reductive monoid with zero ([8],[10]):

Proposition 8. *Let S be a reductive monoid with unit group G . Then there exist a reductive group G_1 and a reductive monoid with zero S_0 such that*

$$S \cong (G_1 \times S_0)/\Gamma,$$

where Γ is a finite central subgroup of G . In particular, if $G_2 = G(S_0)$, then $G \cong (G_1 \times G_2)/\Gamma$. $\square$

It follows that, as in the case of toric varieties, we can restrict ourselves to the study of reductive monoids with zero:

Proposition 9. *Let S be a reductive monoid with unit group G . Let G_1, G_2, S_0 and Γ be as above. Then, under the identification $\mathcal{Q}(G) = \mathcal{Q}(G_2) \oplus \mathcal{Q}(G_1)$, we have:*

$$(3) \quad (\mathcal{C}(S), \mathcal{F}(S)) = (\{0\} \times \mathcal{C}(S_0), \{0\} \times \mathcal{D}(G_1)) .$$

PROOF. By Propositions 8 and 7, we can suppose without loss of generality that $G = G_1 \times G_2$, and $S = G_1 \times S_0$, S_0 a monoid with zero and with unit group G_2 . In particular, $\Xi(G) = \Xi(G_1) \times \Xi(G_2)$, and $\mathcal{Q}(G) = \mathcal{Q}(G_1) \oplus \mathcal{Q}(G_2)$. If $B_i \subset G_i$ is a Borel subgroup of G_i , $i = 1, 2$, then $B = B_1 \times B_2$ is a Borel subgroup of G . If $\alpha_1, \dots, \alpha_r$ are the simple roots associated to B_1 , and $\beta_1, \dots, \beta_s$ those associated to B_2 , then $(\alpha_1, 0), \dots, (\alpha_r, 0), (0, \beta_1), \dots, (0, \beta_s)$ are the simple roots associated to B .

Denote D_{α_i} and D_{β_j} the colours of G_1 and G_2 respectively. It is immediate that the colours of G are:

$$\mathcal{D}(G) = \{D_{\alpha_i} \times G_2, G_1 \times D_{\beta_j} : i = 1, \dots, r, j = 1, \dots, s\} .$$

As $k[G] = k[G_1] \otimes k[G_2]$, the valuations associated to $D_{\alpha_i} \times G_2$ and $G_1 \times D_{\beta_j}$ are $(\alpha_i^\vee, 0)$ and $(0, \beta_j^\vee)$ respectively. In the same way, we see that $\mathcal{V}(G) = -C(G) = -(C(G_1) \times C(G_2))$.

A divisor $D \subset S$ is $(G \times G)$ -stable if and only if $D = G_1 \times D'$, where D' is a divisor $(G_2 \times G_2)$ -stable of S_0 . It follows that the valuation associated to D satisfies $\rho_G(\nu_D) = (0, \rho_{G_2}(\nu_{D'}))$.


On the other hand, the kernel of S (i.e. the unique $(G \times G)$ -closed orbit) is equal to $G_1 \times \{0\}$. It follows that the set of colours $\mathcal{F}(S)$ is $\{0\} \times \mathcal{D}(G_2)$ (we recall that the set of colours of a monoid with zero S_0 is $\mathcal{D}(G(S_0))$), and the proposition is proved. $\square$

Corollary 4. *Let S be a reductive monoid with unit group G . Let $(\mathcal{C}(S), \mathcal{F}(S))$ be the colored cone associated to S . Then $\mathcal{C}(S) + \mathbb{Q}^+ \mathcal{D}$ is a strictly convex cone.* $\square$

The converse to this corollary is not true, as the following example shows:

Example 2. Consider $G = \mathrm{Sl}(2) \times k^*$. We identify $\mathcal{Q}(G)$ to $\mathbb{Q}^2$, and $\Xi_*(G)$ to the lattice $\mathbb{Z}^2 \subset \mathbb{Q}^2$. The affine embeddings of G are exactly those where the closed orbit is a point (with colored cone of the form $(\mathcal{C}, (\alpha^\vee, 0))$), and the *elementary embeddings* associated to the half lines $(\mathbb{Q}^+(0, 1), \emptyset)$ and $(\mathbb{Q}^+(0, -1), \emptyset)$.

Indeed, let $(\mathcal{C}, \emptyset)$ be the colored cone associated to an affine embedding without colors. Then there exists $\chi \in \Lambda$ such that $\chi(\alpha^\vee, 0) > 0$, $\chi(\nu) \leq 0$ for all $\nu \in \mathcal{V}$. It follows that $\chi(\nu) = 0$ if and only if ν belongs to the y -axis. Let $\nu \in \mathcal{V}$ such that the cone $\mathcal{C}'$ generated by ν and $(\alpha^\vee, 0)$ is strictly convex. Then the embedding S associated to $(\mathcal{C}', (\alpha^\vee, 0))$ is affine (just take $\chi \equiv 0$). In particular, S is a reductive monoid (with zero: the set of colors of S is all $\mathcal{D}$).



On the other hand, if we take $\tau = \mathcal{C}' \cap \mathcal{V}$, then $\tau + \mathbb{Q}^+(\alpha^\vee, 0) = \mathcal{C}'$ is strictly convex, but the embedding associated to the colored cone $(\tau, \emptyset)$ is not affine.

An immediate consequence of the preceding proposition is that the colored cone of a reductive monoid S is determined by $\mathcal{V}(S)$, or even $\mathcal{B}(S)$. Moreover, we can find the decomposition of S as a quasi-direct product of a reductive monoid with zero and a reductive group (cf. Proposition 8) as follows:

If we keep the notation of Proposition 8, the space $\mathcal{Q}(G_2)$ is $\langle \mathcal{C}(S) \rangle$, the subspace of $\mathcal{Q}(G)$ generated by $\mathcal{C}(S)$. The lattice considered contains $\Xi(G) \cap \langle \mathcal{C}(S) \rangle$ as a subgroup of finite index.

A simply connected covering of the commutator of the normal subgroup G_1 (resp. G_2) is the semi-simple simply connected group whose root system is given by the Dynkin diagram of G restricted to $\mathcal{F}(S)$ (resp. $\mathcal{D}(G) \setminus \mathcal{F}(S)$).

In terms of the combinatorics of the colored cones of $\mathcal{Q}(G)$, this remark gives the following result:

Proposition 10. *Let G be a reductive group. Given a polyhedral cone τ contained in $\mathcal{V}(G) = -C$, such that $\tau + \mathbb{Q}^+\mathcal{D}$ is strictly convex, there exists a unique subset of colours $\mathcal{F} \subset \mathcal{D}$ such that $(\tau + \mathbb{Q}^+\mathcal{F}, \mathcal{F})$ is the colored cone associated to an affine embedding of G .*

PROOF. Let $\tau \subset \mathcal{V}(G)$ be a cone such that $\tau + \mathbb{Q}^+\mathcal{D}$ is strictly convex. If A is a subset of $\mathcal{Q}(G)$, denote $\langle A \rangle$ the sub-space of $\mathcal{Q}(G)$ generated by A . Consider the projection $p : \mathcal{Q}(G) \rightarrow \mathcal{Q}(G_0)$.

By the preceding remark, $\mathcal{F} = \mathcal{D} \cap \langle p(\tau) \rangle$ is the natural candidate for $(\tau + \mathbb{Q}^+\mathcal{F}, \mathcal{F})$ to be a colored cone corresponding to an affine embedding of G . Let $\{v_1, \dots, v_r\} \subset \mathcal{V}(G)$ be generators of the cone τ . Then $\tau + \mathbb{Q}^+\mathcal{F}$ is generated by $\{v_1, \dots, v_r\} \cup \mathcal{F} \subset \mathcal{V}(G) \cup \mathcal{D}$. The cone $\tau + \mathbb{Q}^+\mathcal{D}$ is strictly convex, and thus the same holds for $\tau + \mathbb{Q}^+\mathcal{F}$.

Let us show that $(\text{int}(\tau + \mathbb{Q}^+\mathcal{F})) \cap \mathcal{V}(G)$ is non-void. Then $(\tau + \mathbb{Q}^+\mathcal{F}, \mathcal{F})$ will be a colored cone.

Let $v = \sum_{n_j > 0} n_j v_j \in \text{int } \tau$. We have:

$$p(v) \in p(\text{int } \tau) \subset p(\mathcal{V}(G)) = -C(G_0).$$

If $q > 0$ is sufficiently small, as the set $\mathcal{F}$ is contained in $\langle p(\tau) \rangle$, we have

$$p \left(v + \sum_{(\alpha^\vee, 0) \in \mathcal{F}} q(\alpha^\vee, 0) \right) = p(v) + \sum_{(\alpha^\vee, 0) \in \mathcal{F}} q(\alpha^\vee, 0) \in \text{int}(p(\tau)).$$

It follows that

$$v + q \sum_{(\alpha^\vee, 0) \in \mathcal{F}} (\alpha^\vee, 0) \in (\text{int}(\tau + \mathbb{Q}^+\mathcal{F})) \cap p^{-1}(-C(G_0)) = (\text{int}(\tau + \mathbb{Q}^+\mathcal{F})) \cap \mathcal{V}(G),$$

and thus $(\tau + \mathbb{Q}^+\mathcal{F}, \mathcal{F})$ is a colored cone.

In order to prove that $(\tau + \mathbb{Q}^+\mathcal{F}, \mathcal{F})$ corresponds to an affine embedding, we consider a basis $\mathcal{F} \cup \{w_1, \dots, w_s\}$ of $\langle \tau \rangle + \langle \mathcal{F} \rangle$, where w_i belongs to $\Xi(G)$ for all $i = 1, \dots, s$. By construction, $(\langle \tau \rangle + \langle \mathcal{F} \rangle) \cap \langle \mathcal{D} \setminus \mathcal{F} \rangle = \{0\}$. We can then add the elements of $\mathcal{D} \setminus \mathcal{F}$ to $\mathcal{F} \cup \{w_1, \dots, w_s\}$ and obtain in this way a linearly independent set. We complete this set in a basis $\mathcal{B}$ of $\mathcal{Q}(G)$:

$$\mathcal{B} = \mathcal{D} \cup \{w_1, \dots, w_s\} \cup \{w_{s+1}, \dots, w_h\} \quad w_i \in \Xi(G), \quad i = 1, \dots, h.$$

Define $\chi \in \Xi_+(G)$ over $\mathcal{B}$ as follows:

$$\chi(b) = \begin{cases} 1 & \text{if } b \in \mathcal{D} \setminus \mathcal{F} \\ 0 & \text{otherwise} \end{cases}$$

and extend χ by linearity. It is easy to see that χ satisfies the conditions a)–c) of Proposition 5 (iv), so the embedding associated to $(\tau + \mathbb{Q}^+\mathcal{F}, \mathcal{F})$ is affine.

It remains to prove the unicity of the chosen set of colours. Let $\mathcal{F}' \subset \mathcal{D}$ be a subset of colours such that $(\tau + \mathbb{Q}^+\mathcal{F}', \mathcal{F}')$ is a colored cone corresponding to an affine embedding of G . Let us show that $\mathcal{F}' = \mathcal{F}$.

Let $v \in (\text{int}(\tau + \mathbb{Q}^+ \mathcal{F}')) \cap \mathcal{V}(G) = (\text{int} \mathcal{C}(S')) \cap \mathcal{V}(G)$. The vector v can be written as

$$v = v_0 + \sum_{(\alpha, 0) \in \mathcal{F}'} q_\alpha(\alpha^\vee, 0) \quad , \quad v_0 \in \tau, \quad q_\alpha > 0 .$$

Let $\chi \in \Lambda$ be as above. As $\chi(w) \leq 0$ for all $w \in \mathcal{V}(G) = -C(G)$, and $\chi > 0$ over $\mathcal{D} \setminus \mathcal{F}'$, it follows that $0 \geq \chi(v) = \sum_{(\alpha, 0) \in \mathcal{F}' \setminus \mathcal{F}} q_\alpha \chi(\alpha^\vee, 0)$. The only possibility is that $\chi(v) = 0$, and $\mathcal{F}' \subset \mathcal{F}$. By symmetry, we get $\mathcal{F} \subset \mathcal{F}'$. $\square$

Corollary 5. *Let G be a reductive group. Let S and S' be two monoids with unit group G , such that $\mathbb{Q}^+ \mathcal{B}(S) = \mathbb{Q}^+ \mathcal{B}(S')$. Then S and S' are isomorphic. In particular, if $\mathcal{V}(S) = \mathcal{V}(S')$, then S and S' are isomorphic.* $\square$

Remark. Let us give a geometric interpretation of the corollary above. If S is a reductive monoid with unit group G and $(\mathcal{C}(F), \mathcal{F}(S))$ its colored cone, there exists a proper G -morphism $\tilde{S} \rightarrow S$, where $\tilde{S}$ is the embedding associated to $(\mathcal{V}(S), \emptyset)$ ([1]). Moreover, the embedding $\tilde{S}$ is minimal for this property; it is called the *decoloration* of S . Corollary 5 says then that two monoids are isomorphic if and only if their decolorations are.

In conclusion, we have proved the following classification:

Theorem 4. *Let G be a reductive group. The isomorphism classes of algebraic monoids with unit group G are in bijection with the strictly convex polyhedral cones of $\mathcal{Q}(G)$ generated by all the colours and a finite set of elements of $\mathcal{V}(G)$.*

5. SOME CONSEQUENCES

We end up with some applications of the classification obtained above.

First, we describe the algebra of regular functions on a reductive monoid, generalising the result obtained by Vinberg in characteristic zero ([11]).

Notation. Let G be a reductive group. We keep the notations of the last section. If M is a G -module, we note $M^{(B)}$ the set of eigenvectors of M for the action of B .

Proposition 11. *Let S be a monoid with unit group G . Then,*

$$k[S \setminus \overline{\cup_{D \in \mathcal{F}(S)} D}]^{(B \times B^-)} = \left\{ f \in k(G)^{(B \times B^-)} : \chi_f \in \mathcal{C}(S)^\vee \right\} , \quad \text{and}$$

$$k[S]^{(B \times B^-)} = \left\{ f \in k(G)^{(B \times B^-)} : \chi_f \in \mathcal{C}(S)^\vee \cap \Xi_+(G) \right\} .$$

Moreover, the set $\mathcal{L}_S = \mathcal{C}(S)^\vee \cap \Xi(T)$ of weights of $k[S]^{(B \times B^-)}$ generates $\Xi(T)$, the set of weights of $k(G)^{(B \times B^-)} = k(S)^{(B \times B^-)}$.

In particular, if $\text{car } k = 0$, the decomposition of $k[S]$ into simple $(G \times G)$ -modules is the following:

$$k[S] = \bigoplus_{\chi \in \mathcal{C}(S)^\vee \cap \Xi_+(G)} V_\chi \otimes V_\chi^* .$$

PROOF. The first equality follows from [5, thm. 3.5]. In order to prove the second one, observe that a function $f \in k[S \setminus \overline{\bigcup_{D \in \mathcal{F}(S)} D}]^{(B \times B^-)}$ extends to a regular function on S if and only if $\nu_D(f) \geq 0$ for all $D \in \mathcal{F}(S)$. By proposition 6, this means that $\langle \chi_f, \alpha_i^\vee \rangle = \nu_{D_i}(f) \geq 0$, i.e. χ_f is a dominant weight.

Let $f = \psi/\varphi \in k(S)^{(B \times B^-)}$, where $\varphi, \psi \in k[S]$. The $(G \times G)$ -module $V \subset k[S]$ generated by φ is of finite dimension. Let $\{g_1 \cdot \varphi, \dots, g_r \cdot \varphi\}$, $g_i \in G \times G$, be a system of generators of V . Then, there exists a $(B \times B^-)$ -eigenvector $h = \sum_{i=1}^r t_i g_i \cdot \varphi \in V$, $t_i \in k$. Consider the $(B \times B^-)$ -eigenvector $hf = (\sum_{i=1}^r t_i g_i \cdot \varphi)\psi/\varphi \in k[S]$; its weight is $\chi_{hf} = \chi_h + \chi_f$. We have $f = hf/h$, where $hf, h \in k[S]^{(B \times B^-)}$, and $\chi_f = \chi_{fh} - \chi_h$. It follows that $\mathcal{L}_S$ generates $\Xi(T)$.

Finally, the last equality is a direct consequence of the complete reducibility of the rational representations of $G \times G$ in characteristic 0. $\square$

By Proposition 4, the ideals of a reductive monoid S are exactly the closed $(G \times G)$ -stable subvarieties of S . The dimension of an irreducible component of an ideal is then given by the dimension of the open orbit in this component. By Proposition 5 (iii), to calculate those dimensions it suffices to calculate the stabilizers of the colours. But the Bruhat decomposition gives us this information immediately:

Proposition 12 ([3, §29]). *Let G be a reductive group. If we keep the preceding notations, the stabilizer in $G \times G$ of the colour $D_i = \overline{Bs_{\alpha_i}B^-}$, $i = 1, \dots, l$, is generated by the parabolic subgroups $(P_{\alpha_j} \times P_{-\alpha_j})$, $j \neq i$, where P_α denotes the minimal parabolic associated to the simple root α .* $\square$

Corollary 6. *Let S be a reductive monoid with unit group G . If $Y \subset S$ is a $(G \times G)$ -orbit, whose combinatorial data is $(\mathcal{C}_Y(S), \mathcal{F}(Y))$, its dimension is given by:*

$$\dim Y = l + \dim Z(G) - \dim \mathcal{C}_Y(S) + \dim G/P_{\mathcal{D} \setminus \mathcal{F}(Y)}$$

where $P_{\mathcal{D} \setminus \mathcal{F}(Y)}$ is the parabolic subgroup generated by the parabolics P_α , $\alpha \in \mathcal{D} \setminus \mathcal{F}(Y)$.

PROOF.

Apply Proposition 5 (iii) and then Lemma 12 in order to calculate the intersection of the parabolic subgroups which stabilize the colours belonging to $\mathcal{D} \setminus \mathcal{F}(Y)$. $\square$

REFERENCES

- [1] M. Brion, *Sur la Géométrie des variétés sphériques*. Comment. Math. Helv. 66 (1991), 237–262.
- [2] R. Hartshorne. *Ample subvarieties of Algebraic Varieties*. Lect. Notes in Math 157. Springer Verlag, New York, 1970.
- [3] J. Humpreys, *Linear Algebraic Groups*. GTM 21, Springer Verlag, New York, 1970.
- [4] G. Kempf, *Algebraic Varieties*. London Math. Soc. Lect. Notes 172, Cambridge Univ. Press, 1993.
- [5] F. Knop, *The Luna-Vust Theory of Spherical Embeddings*. In S. Ramanan *et al*, editors, Proceedings of the Hyderabad Conference on Algebraic Groups, pages 225–249. National Board for Higher Mathematics, Manoj, 1991.
- [6] D. Mumford, J. Fogarty and F. Kirwan, *Geometric Invariant Theory*. Modern Surveys in Math. 34, Springer Verlag, New York, 3rd enlarged ed., 1994.
- [7] T. Oda, *Torus embeddings and applications*. Tata Press, Bombay, 1978.
- [8] M.S. Putcha, *Linear algebraic monoids*. London Math. Soc. Lect. Notes 133, Cambridge Univ. Press, 1988. Notes 133,
- [9] L. Renner, *Algebraic monoids*. PhD. Thesis, Univ. of British Columbia, 1982.
- [10] A. Rittatore, *Monoïdes algébriques et variétés sphériques*. PhD. Thesis, Institut Fourier, Grenoble, France, 1997. <http://www-fourier.ujf-grenoble.fr>
- [11] E.B. Vinberg, *On reductive Algebraic Semigroups*. Amer. Math. Soc. Transl., Serie 2, 169 (1994), 145 – 182. Lie Groups and Lie Algebras. E.B. Dynkin seminar.
- [12] T. Vust, *Plongements d’espaces symétriques algébriques: une classification*. Ann. Scuola Norm. Sup. Pisa, XVII, 2 (1990), 165–194.
- [13] W. Waterhouse, *The unit group of affine algebraic monoids*. Proc. Amer. Math. Soc. 85 (1982) 506 – 508.

ALVARO RITTATORE

Institut Fourier,

Grenoble I.

B.P. 74.

38402 St. Martin d’Hères CEDEX France

e-mail: alvaro@fourier.ujf-grenoble.fr